



**Queensland University of Technology**  
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Lin, R & Liu, Fawang (2007) Fractional High Order Methods For The Non-linear Fractional Ordinary Differential Equation. *Nonlinear Analysis*, 66(4), pp. 856-869.

This file was downloaded from: <http://eprints.qut.edu.au/21852/>

**Notice:** *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1016/j.na.2005.12.027>

# Fractional High Order Methods For the Nonlinear Fractional Ordinary Differential Equation

R. Lin <sup>a,1</sup>, F. Liu <sup>a,b,2</sup>

<sup>a</sup>*School of Mathematical Sciences, Xiamen University, Xiamen 361005, China*

<sup>b</sup>*School of Mathematical Sciences, Queensland University of Technology, GPO Box  
2434, Brisbane, Qld. 4001, Australia.*

---

## Abstract

In this paper, we consider the nonlinear fractional order ordinary differential equa-

tion  ${}_0D_t^\alpha y(t) = f(y, t)$ , ( $t > 0$ ),  $n - 1 < \alpha \leq n$ ,  $y^{(i)}(0) = y_0^{(i)}$ ,  $i = 0, 1, 2, \dots, n - 1$ , where  $f(y, t)$  satisfies the  $L$ -condition, i.e.,  $|f(y_1, t) - f(y_2, t)| \leq L|y_1 - y_2|$  in  $t \in [0, T]$ . Fractional order linear multiple step methods are introduced. The high order (2-6) approximations of the fractional order ordinary differential equation with initial value are proposed. The consistence, convergence and stability of the fractional high order methods are proved. Finally, some numerical examples are provided to show that the fractional high order methods for solving the fractional order nonlinear ordinary differential equation are computationally efficient solution methods.

*Key words:* fractional high order methods; nonlinear fractional order; consistence; convergence; stability

---

## 1 Introduction

Various fields of science and engineering deal with dynamical systems, which can be described by fractional-order equations. This topic has received a great

---

<sup>1</sup> *Email address* :3425linran@sohu.com

<sup>2</sup> Corresponding author. *Email address* :fwliu@xmu.edu.cn, f.liu@qut.edu.au

deal of attention in the last decade [7,15,22–24], especially in the fields of viscoelastic materials [1,10,11,27], electrochemical processes [8], dielectric polarization [28], colored noise [29], anomalous diffusion, signal processing [21], control theory [24], advection and dispersion of solutes in natural porous or fractured media [2,3] and chaos [20]. Djrbasjan et al. [6] considered the Cauchy problem with multi-term fractional derivatives, and proved the Cauchy problem has a unique solution. Kilbas et al. [9] gave solution of Volterra integrodifferential equations with generalized Mittag-Leffler function in the kernels. The main reason for the success of the theory in these cases is that these new fractional-order models are more accurate than integer-order models.

Numerical methods associated with integral order ordinary differential equation, for example, Euler method, linear multiple step method, and so on, have been treated extensively in the literature. On the other hand, theoretical studies of the numerical methods and the error estimate of fractional order differential equation are quite limited, because theoretical analysis of fractional order numerical methods is very difficult [13,14,16–18,26].

Recently linear FDEs based on the Reimann-Liouville fractional derivative with general variable coefficients are solved by adapting the decomposition method. Not much has been done for the nonlinear FDE. For example, in the modelling of viscoelasticity, the nonlinear FDE can describe the strain and entropy reacting in response to changes in stress and temperature. In the theory of viscoelasticity, it is well known that the constitutive equations governing these phenomena involve differential equations of fractional order. Since the theory of viscoelasticity is essentially a linear theory, these differential equations are also linear, and therefore they may be solved using rather simple methods. We shall see that, in our situation, we have to replace these linear equations by nonlinear ones. The standard solution methods for the linear equations usually fail in the nonlinear case. Thus we have also developed an algorithm for the numerical solution of our equations.

There have been some attempts to solve linear problems, but a complete analysis has not been given so far [25]. Nonlinear equations have received rather less attention in the literature, partly because many of the model equations proposed have been linearized. More recently, applications have included classes of nonlinear fractional differential equations and this motivates us to consider their effective numerical methods for the solution of nonlinear fractional differential equations. Diethelm et al. have done a lot of excellent works on numerical methods for fractional order ordinary differential equations [4,5]. In this paper, we propose fractional high order numerical methods for the nonlinear fractional ordinary differential equation.

This paper is organized as follows: Section 2 presents fractional high order numerical methods. In section 3, we introduce the notations, definitions, and

lemmas which we shall use. In sections 4 to 6, we analyze the error theory of the fractional high order numerical methods for the nonlinear fractional differential equation. The consistence, convergence and stability of the fractional high order numerical methods are proven. In section 7, we give some numerical examples based on the use of the high order (2-6) approximations for solving the nonlinear fractional order ordinary differential equation. We show that the high order approximations are computationally efficient methods. Finally, we give a conclusion based on the algorithm for solving the nonlinear fractional order ordinary differential equation.

## 2 High order fractional linear multiple step methods

In this paper, we consider the following nonlinear fractional order ordinary equation (NFOODE)

$${}_0D_t^\alpha y(t) = f(y, t), \quad (t > 0), y^{(i)}(0) = y_0^{(i)}, \quad i = 0, 1, \dots, n-1, \quad (1)$$

where  $n-1 < \alpha < n$ .

In Eq. (1),  $f(y, t)$  satisfies the  $L$ -condition, i.e.,  $|f(y_1, t) - f(y_2, t)| \leq L|y_1 - y_2|$  in  $t \in [0, T]$ . Eq. (1) uses Riemann-Liouville derivative of order  $\alpha$ , which is defined by

$${}_0D_t^\alpha f(t) = \frac{d^n}{dt^n} \left( \frac{1}{\Gamma(n-\alpha)} \int_0^t \frac{f(\tau) d\tau}{(t-\tau)^{\alpha-n+1}} \right).$$

Using the relationship between Grünwald-Letnikov and Riemann-Liouville fractional derivatives ([24]), we have

$${}_0D_t^\alpha f(t) = \lim_{h \rightarrow 0} h^{-\alpha} \Delta_h^\alpha f(t) = \lim_{h \rightarrow 0} h^{-\alpha} \sum_{j=0}^{\lfloor t/h \rfloor} (-1)^j \binom{\alpha}{j} f(t-jh),$$

where  $0 \leq n-1 < \alpha \leq n$ . *Lubich* has given the  $p$ -th ( $p = 1, 2, 3, 4, 5$ , and  $6$ ) approximations [19]

$${}_0D_t^\alpha y(t) \approx {}_0\widetilde{D}_t^\alpha y(t) = h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y(t_{n-j}) + h^{-\alpha} \sum_{j=0}^s w_{nj} y(t_j), \quad nh = T. \quad (2)$$

So we obtain the approximations of order 1, 2, 3, 4, 5, and 6 in the form of (2), where the coefficients  $\omega_k^{(\alpha)}$  are the coefficients of the Taylor series expansions of the corresponding generating functions  $W_p^{(\alpha)}(x)$ , where  $p$  denote the order of method. When  $p$  changes from 1 to 6, the corresponding  $W_p^{(\alpha)}(x)$  are as follows:

$$W_1^{(\alpha)}(x) = (1-x)^\alpha,$$

$$\begin{aligned}
W_2^{(\alpha)}(x) &= \left(\frac{3}{2} - 2x + \frac{1}{2}x^2\right)^\alpha, \\
W_3^{(\alpha)}(x) &= \left(\frac{11}{6} - 3x + \frac{2}{3}x^2 - \frac{1}{3}x^3\right)^\alpha, \\
W_4^{(\alpha)}(x) &= \left(\frac{25}{12} - 4x + 3x^2 - \frac{1}{3}x^3 + \frac{1}{4}x^4\right)^\alpha, \\
W_5^{(\alpha)}(x) &= \left(\frac{137}{60} - 5x + 5x^2 - \frac{10}{3}x^3 + \frac{5}{4}x^4 - \frac{1}{5}x^5\right)^\alpha, \\
W_6^{(\alpha)}(x) &= \left(\frac{147}{60} - 6x + \frac{15}{2}x^2 - \frac{20}{3}x^3 + \frac{15}{4}x^4 - \frac{6}{5}x^5 + \frac{1}{6}x^6\right)^\alpha.
\end{aligned}$$

And the starting weights  $w_{nj}(j = 1, \dots, s)$  are independent of  $h$ . It can be obtained from system of linear equations,

$$\sum_{j=1}^s w_{nj} j^q = \frac{\Gamma(q+1)}{\Gamma(-\alpha+q+1)} n^{q-\alpha} - \sum_{j=1}^s \omega_{n-j}^{(\alpha)} j^q, \quad (q = 0, \dots, s-1),$$

where we can take  $s = p$  under certain condition ([19]).

Using the relationship between Grünwald-Letnikov and Riemann-Liouville fractional derivatives, and above discrete approximations to fractional derivative term, we obtain  $p$ -order (high order,  $p \geq 1$ ) fractional linear multiple step methods ( $p$ -HOFLMSM):

$$\begin{aligned}
h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} y_j &= f(y_n, t_n), \\
y^{(i)}(0) &= y_0^{(i)}, \quad i = 0, 1, \dots, n-1,
\end{aligned} \tag{3}$$

where  $0 < n-1 < \alpha \leq n$ .

### 3 Preparation

To motivate the following definitions, firstly we consider the case  $\alpha = 1$  in (1). If the  $p$ -HOFLMSM ( $p = 1$ ) is applied to the first order ordinary differential equation

$$y'(t) = f(t, y), \quad y(0) = y_0.$$

It is well known that the resulting numerical solution can be written as the Euler method where the weight  $\omega_j^1, j=1, 2, \dots$ , are the coefficients of

$$W_1^{(1)}(x) = 1 - x.$$

Similar to the consistence and stability of the Euler method, we can define them for  $p$ -HOFLMSM.

**Definition 3.1**  $p$ -HOFLMSM is consistent if  $T_n(t) = y(t_n) - \widetilde{y}_n \rightarrow 0$ , as  $h \rightarrow 0$ , where  $\widetilde{y}$  is a solution of  $p$ -HOFLMSM (3) and  $y(t)$  is a solution of NFOODE (1).

**Definition 3.2**  $p$ -HOFLMSM is stable if any two solutions  $y_n$  and  $y'_n$  of  $p$ -HOFLMSM (3), which use different initial values  $f$  and  $f'$ , satisfy  $|y_n - y'_n| \leq C\|f - f'\|_\infty$ , where  $C$  is a constant.

**Lemma 3.3** Suppose that  $f(t)$  is sufficiently differentiable in  $[a, b]$ . Then

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} f(t_{n-j}) + h^{-\alpha} \sum_{j=0}^s w_{nj} f(t_{n-j}) - {}_0D_t^\alpha f(t) = O(h^p),$$

uniformly in  $t \in [a, b]$ , and  $\omega_j^{(\alpha)} = O(j^{-\alpha-1})$ .

**Proof.** ([19] p.707).

**Lemma 3.4** Let  $\omega_k^{(\alpha)} (k = 0, 1, 2, \dots)$  is the coefficient of the Taylor series expansions of the corresponding "generating" functions,  $W_p^{(\alpha)}(x)$ ,  $d_k (k = 0, 1, 2, \dots)$  is the coefficient of  $1/W_p^{(\alpha)}(x)$ , and

$$A = \begin{pmatrix} \omega_0^{(\alpha)} & \omega_1^{(\alpha)} & \dots & \omega_n^{(\alpha)} \\ 0 & \omega_0^{(\alpha)} & \dots & \omega_{n-1}^{(\alpha)} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \omega_0^{(\alpha)} \end{pmatrix}, \quad D = \begin{pmatrix} d_0 & d_1 & \dots & d_n \\ 0 & d_0 & \dots & d_{n-1} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & d_0 \end{pmatrix}.$$

We have  $A^{-1} = D$  and  $d_k = O(k^{\alpha-1})$ .

**Proof.** Let  $AD = B$ . Easily we have  $b_{i,i} = \omega_0^{(\alpha)} d_0, i = 0, \dots, n$  and  $b_{i,j} = 0, i > j$ . And if  $i < j$ ,  $b_{i,j} = \omega_0^{(\alpha)} d_{j-i} + \dots + \omega_{j-i}^{(\alpha)} d_0$ . Let  $f(x) = W_p^{(\alpha)}(x)/W_p^{(\alpha)}(x)$ . It implies that

$$\begin{aligned} f(x) &= (\omega_0^{(\alpha)} + \omega_1^{(\alpha)} x + \dots) \cdot (d_0 + d_1 x + \dots) \\ &= \omega_0^{(\alpha)} d_0 + (\omega_1^{(\alpha)} d_0 + \omega_0^{(\alpha)} d_1) x + \dots + (\omega_k^{(\alpha)} d_0 + \dots + \omega_0^{(\alpha)} d_k) x^k + \dots \end{aligned}$$

From  $f(x) = 1$ , we have  $\omega_0^{(\alpha)} d_0 = 1$ , and if  $i < j$ ,  $b_{i,j} = 0$ . Hence,  $AD = B = E$ . And from [19], we have  $d_k = O(k^{\alpha-1})$ .

**Lemma 3.5** (Gronwall inequality) *Let  $a, b \geq 0$ , and  $\{\eta_i\}$  satisfy*

$$|\eta_n| \leq b + ah \sum_{i=0}^{n-1} |\eta_i|, \quad n = k, k+1, \dots, nh \leq T,$$

*then*

$$|\eta_n| \leq e^{aT} (b + akhM_0), \quad n \geq k, \quad nh \leq T,$$

*where  $M_0 = \max(|\eta_0|, |\eta_1|, \dots, |\eta_{k-1}|)$ .*

**Proof.**([12])

Looking at the questions of existence and uniqueness of the NFOODE, it can be obtained from the following results that are very similar to the corresponding classical theorems in the case of first-order ordinary differential equation. The following theorems have been proven by Diethelm et al. [4].

**Theorem 3.6** (existence) *Assume that  $\Omega = [y_0^{(0)} - \beta, y_0^{(0)} + \beta] \times [0, \chi^*]$  with some  $\chi^* > 0$  and some  $\beta > 0$ , and let the function  $f : \Omega \rightarrow R$  be continuous. Furthermore, define  $\chi = \min\{\chi^*, \alpha\Gamma(q+1)/\|f\|_\infty\}$ . Then, there exists a function  $y : [0, \chi] \rightarrow R$  solving the initial value problem (1).*

**Theorem 3.7** (uniqueness) *Assume that  $\Omega = [y_0^{(0)} - \beta, y_0^{(0)} + \beta] \times [0, \chi^*]$  with some  $\chi^* > 0$  and some  $\beta > 0$ , and let the function  $f : \Omega \rightarrow R$  be bounded on  $\Omega$  and fulfill a Lipschitz condition with respect to the first variable; i.e.,*

$$|f(y, t) - f(z, t)| \leq L |y - z|,$$

*with some constant  $L > 0$  independent of  $t, y$ , and  $z$ . Then denoting  $\chi$  as in Theorem 3.6, there exists at most one function  $y : [0, \chi] \rightarrow R$  solving the initial value problem (1).*

## 4 Consistence

Now we start to present the main results about the error analysis of the fractional order linear multiple step methods.

We consider the error between exact solution  $y(t_n)$  and numerical solution  $\tilde{y}_n$  given from  $y(t_0), \dots, y(t_{n-1})$ .

**Theorem 4.1**  *$p$ -HOFLMSM (3) for solving NFOODE (1) is consistent, and*

$$\tilde{y}_n - y(t_n) = O(h^{p+\alpha}).$$

**Proof.** Let  $n > s$ . We have

$$h^{-\alpha} \omega_0^{(\alpha)} \tilde{y}_n + h^{-\alpha} \sum_{j=1}^n \omega_j^{(\alpha)} y(t_{n-j}) + h^{-\alpha} \sum_{j=0}^s w_{nj} y(t_j) = f(\tilde{y}_n, t_n),$$

$$h^{-\alpha} \omega_0^{(\alpha)} \tilde{y}_n - h^{-\alpha} \omega_0^{(\alpha)} y(t_n) + h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y(t_{n-j}) + h^{-\alpha} \sum_{j=0}^s w_{nj} y(t_j) = f(\tilde{y}_n, t_n),$$

$$h^{-\alpha} \omega_0^{(\alpha)} (\tilde{y}_n - y(t_n)) + {}_0D_t^\alpha y(t) - C_n h^p = f(y(t_n), t_n) + f(\tilde{y}_n, t_n) - f(y(t_n), t_n),$$

$$h^{-\alpha} \omega_0^{(\alpha)} (\tilde{y}_n - y(t_n)) = C_n h^p + (f(\tilde{y}_n, t_n) - f(y(t_n), t_n)),$$

$$(\tilde{y}_n - y(t_n)) = C_n h^{p+\alpha} + \frac{h^\alpha}{\omega_0^{(\alpha)}} (f(\tilde{y}_n, t_n) - f(y(t_n), t_n)),$$

$$\begin{aligned} |\tilde{y}_n - y(t_n)| &\leq C_n h^{p+\alpha} + \left| \frac{h^\alpha}{\omega_0^{(\alpha)}} (f(\tilde{y}_n, t_n) - f(y(t_n), t_n)) \right| \\ &\leq C_n h^{p+\alpha} + \frac{L h^\alpha}{\omega_0^{(\alpha)}} |(\tilde{y}_n - y(t_n))|, \end{aligned}$$

$$\left(1 - \frac{L h^\alpha}{\omega_0^{(\alpha)}}\right) |(\tilde{y}_n - y(t_n))| \leq C_n h^{p+\alpha}.$$

As  $h$  is sufficiently small,  $|(\tilde{y}_n - y(t_n))| \leq C h^{p+\alpha}$ .

## 5 Convergence

**Theorem 5.1** *The  $p$ -HOFLMSM (3) to NFOODE (1) is convergent, and  $|y_n - y(t_n)| = O(h^p)$ .*

**Proof.** Let  $e_n = y(t_n) - y_n$ , and we assume  $e_0, e_1, \dots, e_s$  equal to zero.

From

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} y_j = f(y_n, t_n)$$

and

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y(t_{n-j}) + h^{-\alpha} \sum_{j=0}^s w_{nj} y(t_j) = C_n h^p + f(y(t_n), t_n),$$

we get

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} e_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} e_j = C_n h^p + f(y(t_n), t_n) - f(y_n, t_n).$$



Let

$$f(y(t_n), t_n) - f(y_n, t_n) = L_n(y(t_n) - y_n) = L_n e_n.$$

Let  $|L_n| \leq L$  and  $|C_n| \leq C$  for any  $n$ . Now we estimate  $e_n$ . We can see that

$$\begin{aligned} & (h^{-\alpha} \omega_0^{(\alpha)} - L_n) e_n + h^{-\alpha} \omega_1^{(\alpha)} e_{n-1} + \cdots + h^{-\alpha} \omega_{n-(s+1)}^{(\alpha)} e_{s+1} \\ &= C_n h^p - h^{-\alpha} \sum_{j=0}^s (w_{nj} + \omega_{n-j}^{(\alpha)}) e_j, \\ & (h^{-\alpha} \omega_0^{(\alpha)} - L_{n-1}) e_{n-1} + h^{-\alpha} \omega_1^{(\alpha)} e_{n-2} + \cdots + h^{-\alpha} \omega_{n-(s+2)}^{(\alpha)} e_{s+1} \\ &= C_{n-1} h^p - h^{-\alpha} \sum_{j=0}^s (w_{n-1,j} + \omega_{n-1-j}^{(\alpha)}) e_j, \\ & \quad \dots\dots\dots \\ & (h^{-\alpha} \omega_0^{(\alpha)} - L_{s+1}) e_{s+1} = C_{s+1} h^p - h^{-\alpha} \sum_{j=0}^s (w_{s+1,j} + \omega_{s+1-j}^{(\alpha)}) e_j. \end{aligned}$$

Let

$$\begin{aligned} A &= \begin{pmatrix} \omega_0^{(\alpha)} & \omega_1^{(\alpha)} & \cdots & \omega_{n-(s+1)}^{(\alpha)} \\ 0 & \omega_0^{(\alpha)} & \cdots & \omega_{n-(s+2)}^{(\alpha)} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & \omega_0^{(\alpha)} \end{pmatrix}, \\ \vec{e} &= \begin{pmatrix} e_n \\ e_{n-1} \\ \cdots \\ e_{s+1} \end{pmatrix}, \vec{b} = \begin{pmatrix} b_n \\ b_{n-1} \\ \cdots \\ b_{s+1} \end{pmatrix}, B = \begin{pmatrix} L_n & 0 & \cdots & 0 \\ 0 & L_{n-1} & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & L_{s+1} \end{pmatrix}, \\ A^{-1} = D &= \begin{pmatrix} d_0 & d_1 & \cdots & d_{n-(s+1)} \\ 0 & d_0 & \cdots & d_{n-(s+2)} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & d_0 \end{pmatrix}. \end{aligned}$$

Write  $h^{-\alpha} A \vec{e} - B \vec{e} = \vec{b}$ . We have

$$\vec{e} - h^\alpha D B \vec{e} = h^\alpha D \vec{b},$$

$$\begin{aligned}
& \begin{pmatrix} e_n \\ e_{n-1} \\ \dots \\ e_{s+1} \end{pmatrix} - \begin{pmatrix} L_n h^{(\alpha)} d_0 & L_{n-1} h^{(\alpha)} d_1 & \dots & L_{s+1} h^{(\alpha)} d_{n-(s+1)} \\ 0 & L_{n-1} h^{(\alpha)} d_0 & \dots & L_{s+2} h^{(\alpha)} d_{n-(s+2)} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & L_{n-(s+1)} h^{(\alpha)} d_0 \end{pmatrix} \begin{pmatrix} e_n \\ e_{n-1} \\ \dots \\ e_{s+1} \end{pmatrix} \\
&= \begin{pmatrix} h^{(\alpha)} d_0 & h^{(\alpha)} d_1 & \dots & h^{(\alpha)} d_{n-(s+1)} \\ 0 & h^{(\alpha)} d_0 & \dots & h^{(\alpha)} d_{n-(s+2)} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & h^{(\alpha)} d_0 \end{pmatrix} \begin{pmatrix} b_n \\ b_{n-1} \\ \dots \\ b_{s+1} \end{pmatrix}.
\end{aligned}$$

Consider the first term of  $\vec{e}$  and  $\vec{b}$ . We obtain

$$\begin{aligned}
& e_n - (L_n h^\alpha d_0, L_{n-1} h^\alpha d_1, \dots, L_{s+1} h^\alpha d_{n-(s+1)}) \begin{pmatrix} e_n \\ e_{n-1} \\ \dots \\ e_{s+1} \end{pmatrix} \\
&= (h^\alpha d_0, h^\alpha d_1, \dots, h^\alpha d_{n-(s+1)}) \begin{pmatrix} b_n \\ b_{n-1} \\ \dots \\ b_{s+1} \end{pmatrix},
\end{aligned}$$

$$(1 - L_n h^\alpha d_0) e_n = \sum_{i=0}^{n-(s+1)} h^\alpha d_i b_{n-i} + \sum_{i=1}^{n-(s+1)} L_{n-i} h^\alpha d_i e_{n-i}.$$

When  $\alpha > 1$ ,

$$|h^{\alpha-1} d_i| \leq C |n^{1-\alpha} i^{\alpha-1}| \leq C,$$

so

$$\left| \sum_{i=1}^n L_{n-1} h^\alpha d_i e_{n-i} \right| \leq C h \sum_{i=1}^n |L_{n-i} e_{n-i}|.$$

And because  $d_i = O(i^{\alpha-1})$ ,

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n |h^\alpha d_i| \leq \lim_{n \rightarrow \infty} \sum_{i=1}^n C' \frac{i^{\alpha-1}}{n^\alpha} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n C' \left(\frac{i}{n}\right)^{\alpha-1} = C' \int_0^1 x^{\alpha-1} dx = C,$$

then

$$|\sum_{i=1}^n h^\alpha d_i b_{n-i}| \leq C \sup(|b_i|) \leq Ch^p.$$

So we have

$$\begin{aligned} |(1 - L_n h^\alpha d_0) e_n| &\leq Ch^p + \sum_{i=1}^{n-(s+1)} |L_{n-i} h^\alpha d_i e_{n-i}| \\ &\leq Ch^p + h \sum_{i=1}^{n-(s+1)} |L_{n-i} h^{\alpha-1} d_i e_{n-i}| \\ &\leq Ch^p + hLC \sum_{i=1}^{n-(s+1)} |e_{n-i}| \\ &\leq Ch^p + hLC \sum_{i=s+1}^{n-1} |e_i|. \end{aligned}$$

As  $h$  is sufficiently small,  $|e_n| \leq Ch^p + hLC \sum_{i=s+1}^{n-1} |e_i|$ . Using *Gronwall* inequality, we can prove that  $e_n = O(h^p)$ .

## 6 Stability

**Theorem 6.1** *Let  $y_n$  and  $y'_n$  are numerical solutions, which the right hand sides are given by  $f$  and  $f'$ , respectively. Then  $|y_n - y'_n| \leq C \|f - f'\|_\infty$  for any  $n$ , i.e., the  $p$ -HOFLLSM (3) to the NFOODE (1) is stable for the term of the right hand side.*

**Proof.** Let  $y_n - y'_n = \varepsilon_n$ .

From

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} y_j = f(y_n, t_n)$$

and

$$h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} y'_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} y'_j = f'(y'_n, t_n),$$

we have

$$\begin{aligned} &h^{-\alpha} \sum_{j=0}^n \omega_j^{(\alpha)} \varepsilon_{n-j} + h^{-\alpha} \sum_{j=0}^s w_{nj} \varepsilon_{n-j} \\ &= f(y_n, t_n) - f(y'_n, t_n) + f(y'_n, t_n) - f'(y'_n, t_n). \end{aligned}$$

Let

$$f(y_n, t_n) - f(y'_n, t_n) = L_n(y_n - y'_n) = L_n \varepsilon_n,$$

$$f(y'_n, t_n) - f'(y'_n, t_n) = \Delta f_n,$$

and

$$|f(y, t) - f'(y, t)| \leq \Delta f.$$

Obviously  $|L_n| \leq L$  for any  $n$ .

Now we estimate  $\varepsilon_n$ , we can see that

$$\begin{pmatrix} h^{(-\alpha)}\omega_0^{(\alpha)} - L_n & h^{(-\alpha)}\omega_1^{(\alpha)} & \cdots & h^{(-\alpha)}\omega_{n-(s+1)}^{(\alpha)} \\ 0 & h^{(-\alpha)}\omega_0^{(\alpha)} - L_{n-1} & \cdots & h^{(-\alpha)}\omega_{n-(s+2)}^{(\alpha)} \\ \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & h^{(-\alpha)}\omega_0^{(\alpha)} - L_{s+1} \end{pmatrix} \begin{pmatrix} \varepsilon_n \\ \varepsilon_{n-1} \\ \cdots \\ \varepsilon_{s+1} \end{pmatrix} \\ = \begin{pmatrix} \Delta f_n \\ \Delta f_{n-1} \\ \cdots \\ \Delta f_{s+1} \end{pmatrix}.$$

Note that  $h^{-\alpha}A - B\vec{\varepsilon} = \vec{b}$ . From the proof above, we easily have

$$(1 - L_n h^\alpha d_0) \varepsilon_n = \sum_{i=0}^{n-(s+1)} h^\alpha d_i b_{n-i} + \sum_{i=1}^{n-(s+1)} L_{n-i} h^\alpha d_i \varepsilon_{n-i}.$$

When  $\alpha > 1$ ,

$$|(1 - L_n h^\alpha d_0) \varepsilon_n| \leq \left| \sum_{i=0}^{n-(s+1)} h^\alpha d_i b_{n-i} \right| + \sum_{i=1}^{n-(s+1)} |L_{n-i} h^\alpha d_i \varepsilon_{n-i}|,$$

we have

$$|(1 - L_n h^\alpha d_0) \varepsilon_n| \leq C \Delta f + \sum_{i=1}^{n-(s+1)} |L_{n-i} h^\alpha d_i \varepsilon_{n-i}| \leq C \Delta f + hCL \sum_{i=s+1}^{n-1} |\varepsilon_i|.$$

Hence,  $|\varepsilon_n| \leq C \Delta f + hCL \sum_{i=s+1}^{n-1} |\varepsilon_i|$ .

Using *Gronwall* inequality, we have  $|\varepsilon_n| \leq C \Delta f + Ch |\varepsilon_{s+1}| \leq \bar{C} \Delta f$ .

Therefore, for the  $p$ -HOFLMSM, if there is a perturbation in right hand side  $f(y, t)$ , the small change would not cause the large error in the numerical solution. Thus the  $p$ -HOFLMSM is stable on right hand term  $f(y, t)$ .

## 7 Numerical examples

In this section we consider an initial value problem for one of the fractional order differential equations appearing in applied problems [24]. We would like to present some numerical examples to show the effect of the  $p$ -HOFLMSM.

Example 1: Fractional oscillation equation with  $\alpha = 1.5$ :

$${}_0D_t^{\frac{3}{2}}y(t) + y(t) = te^{-t}, y^{(k)}(0) = 0, (k = 0, 1).$$

The exact solution is

$$y(t) = \int_0^t G(t-x)xe^{-x}dx, \quad G(t) = t^{\alpha-1}E_{\alpha,\alpha}(-t^{\alpha}),$$

where  $E_{\alpha,\beta}(z)$  is the so-called *Mittag-Leffler* function

$$E_{\alpha,\beta}(z) = \sum_{j=0}^{\infty} \frac{z^j}{\Gamma(\alpha j + \beta)}, \alpha > 0, \beta > 0.$$

A comparison of the  $p$ -HOFLMSM ( $p = 3$ ) and the exact solution is listed in Table 1 and is shown in Figure 1. From Table 1 and Figure 1, it can be seen that our numerical method (3-HOFLMSM) is in excellent agreement with the exact solution and the error between  $p$ -HOFLMSM and exact solution is  $O(h^p)$ .




Fig. 1. A comparison of the  $p$ -HOFLMSM ( $p = 3$ ) and exact solution in Example 1.

Example 2: Fractional oscillation equation with  $1 < \alpha \leq 2$ .

Table 1

Numerical solution and error ( $n = 100, h = 0.2$ )

$t$	3-HOFLMSM	Exact solution	Error
0.8	9.52875892E-02	9.92045725E-02	3.91698327E-03
2.4	3.84651536E-01	3.84983333E-01	3.31797377E-04
4.0	2.01589390E-01	2.01226046E-01	3.63344850E-04
5.6	-1.20968485E-02	-1.23575173E-02	2.60668883E-04
7.2	-4.09010633E-02	-4.09463462E-02	4.52828845E-05
8.8	-7.20316073E-03	-7.18831888E-03	1.48418475E-05
10.4	4.52776157E-03	4.50733840E-03	2.04231665E-05
12.0	4.73026337E-04	4.27574713E-04	4.54516234E-05
13.6	-1.99206193E-03	-2.03395168E-03	4.18897439E-05
15.2	-1.22253777E-03	-1.25642181E-03	3.38840349E-05
16.8	-3.70895123E-04	-4.03356068E-04	3.24609452E-05
18.4	-2.36528966E-04	-2.70331545E-04	3.38025783E-05
20.0	-2.84607314E-04	-3.18767867E-04	3.41605531E-05
The maximum error is $4.13691139E - 03$ in $t = 0.6$			

We are interested in the behaviors of the solution for the fractional oscillation equation:

$${}_0D_t^\alpha y(t) + y(t) = te^{-t}, \quad 1 < \alpha \leq 2, \quad y^{(i)}(0) = 0, \quad i = 0, 1.$$

We show the behaviors of the solutions with  $1 < \alpha \leq 1.5$  and  $1.5 < \alpha \leq 2$  in Figures 2 and 3, respectively.

Example 3 A nonlinear fractional order ordinary differential equation with  $\alpha = 1.5$ :

$${}_0D_t^{\frac{3}{2}} y(t) + y^2(t) = f(t), \quad y^{(i)}(0) = 0, \quad i = 0, 1$$

where

$$f(t) = \frac{\Gamma(6)}{\Gamma(6-\alpha)} t^{5-\alpha} - \frac{3\Gamma(5)}{\Gamma(5-\alpha)} t^{4-\alpha} + \frac{2\Gamma(4)}{\Gamma(4-\alpha)} t^{3-\alpha} + (t^5 - 3t^4 + 2t^3)^2.$$

The exact solution is  $y(t) = t^5 - 3t^4 + 2t^3$ .

We apply the  $p$ -HOFLMSM with step sizes  $h = 0.02$  and  $p = 3$  to obtain the approximation. A comparison of the  $p$ -HOFLMSM ( $p = 3$ ) and the exact solution is listed in Table 2 and is shown in Figure 4. From Table 2 and Figure 4, it can be seen that our numerical method (3-HOFLMSM) is in excellent




Fig. 2. Behaviors of solution with  $1 \leq \alpha \leq 1.5$ .




Fig. 3. Behaviors of solution with  $1.5 \leq \alpha \leq 2.0$ .

agreement with the exact solution and the error between  $p$ -HOFLMSM and exact solution is  $O(h^p)$ .

Example 4 A nonlinear fractional order ordinary differential equation with  $1 < \alpha \leq 2$ :

$${}_0D_t^\alpha y(t) + y^2(t) = f(t),$$


where  $f(t) = e^{-t} \sin(t)$ ,  $t \in [0, 6]$ .

Figure 5 show the behaviors of the nonlinear solution with  $1 < \alpha \leq 2$ .

Table 2

Numerical solution and error ( $n = 100, h = 0.02$ )

$t$	Numerical solution	Exact solution	Error
0.2	1.15167781E-02	1.15200000E-02	-3.22187462E-06
0.4	6.14108077E-02	6.14399999E-02	-2.91922377E-05
0.6	0.12092028	0.12096000	-3.97172672E-05
0.8	0.12284510	0.12288000	-3.48980138E-05
1.0	-1.54449952E-05	0.00000000	-1.54449952E-05
1.2	-0.27646186	-0.27648000	1.81399204E-05
1.4	-0.65849166	-0.65856000	6.83334800E-05
1.6	-0.98289629	-0.98304000	1.43708888E-04
1.8	-0.93286240	-0.93311999	2.57594614E-04
2.0	4.05999983E-04	0.00000000	4.05999983E-04

The maximum error is  $4.05999983E - 04$  in  $t = 2.0$ Fig. 4. A comparison of the  $p$ -HOFLMSM ( $p = 3$ ) and exact solution in Example 3

## 8 Conclusion

Fractional high order methods for a fractional order nonlinear ordinary differential equation have been described. The consistence, convergence and stability are proved. Numerical examples are presented.  $p$ -HOFLMSM for solving the fractional order ordinary differential equation is an effective method.






Fig. 5. behaviors of the nonlinear solution with  $1 < \alpha \leq 2$

**Acknowledgement** The authors gratefully acknowledge the support of the National Natural Science Foundation of China under Grant 10271098.

## References

- [1] R.L. Bagley and R.A. Calico, Fractional order state equations for the control of viscoelastic structures, *Journal of Guidance, Control, and Dynamics* 14(2), (1999).
- [2] D.A. Benson, S.W. Wheatcraft and M.M. Meerschaert, Application of a fractional advection-dispersion equation, *Water Resour. Res.* 36(6) (2000a) 1403-1412.
- [3] D.A. Benson, S.W. Wheatcraft and M.M. Meerschaert, The fractional-order governing equation of Levy motion, *Water Resour. Res.* 36(6) (2000b) 1413-1423.
- [4] K. Diethelm and N.J. Ford, Analysis of fractional differential equations, *Journal of Mathematical Analysis and Applications* 265, (2002), 220-248.
- [5] K. Diethelm, N.J. Ford and A.D. Freed, Details error analysis for a fractional Adams method, *Numerical Algorithms*, 36, (2004), 31-52.
- [6] M.M. Djrbashian and A.B. Nersesian, Fractional derivatives and the Cauchy problem for differential equations of fractional order. *Izv. Acad. Nauk Armjanskoy SSR*, 3(1):3-29, 1968, In Russian.
- [7] F. Huang and F. Liu, The time fractional diffusion and advection-dispersion equation, *ANZIAM J.*, 46, (2005), 317-330.

- [8] M. Ichise, Y. Nagayanagi and T. Kojima, An analog simulation of non-integer order transfer functions for analysis of electrode processes, *Journal of Electroanalytical Chemistry Interfacial Electrochemisry* 33, (1971), 253-265.
- [9] A.A. Kilbas, M. Saigo and R.K. Saxena, Solutions of Volterra integro-differential equations with generalised Mittag-Leffler function in the kernels, *Journal of Integral Equations*, vol 14, N4, (2002), 377-396.
- [10] R.C. Koeller, Polynomial operators, Stieltjes convolution, and fractional colculus in hereditary methabics, *Acta Mechanica* 58, (1984), 299-307.
- [11] R.C. Koeller, Application of fractional calculus to the theory of viscoelasticity, *Journal of Applied Mechanics* 51, (1984), 299-307.
- [12] R. Li, Numerical solution of differential equation, High Education Press, Beijing, China, (1996).
- [13] R. Lin and F. Liu, High order approximations for the fractional ordinary differential equation with initial value problem, *Journal of Xiamen University (Natural Science)*, Vol. 43, No.1, (2004), 25-30.
- [14] R. Lin and F. Liu, Analysis of fractional-order numerical method for the fractional relaxation equation, *Computational Mechanics (CD-ROM)*, ID-362, (2004).
- [15] F. Liu, V. Anh, I. Turner and P. Zhuang, Time fractional advection-dispersion equation, *J. Appl. Math. Comp.*, (2003), 233-246.
- [16] F. Liu, V. Anh and I. Turner, Numerical solution of space fractional Fokker-Planck equation, *J. Comp. and Appl. Math.*, 166, (2004), 209-219.
- [17] F. Liu, V. Anh, I. Turner and P. Zhuang, Numerical simulation for solute transport in fractal porous media, *ANZIAM J.* 45(E), (2004), 461-473.
- [18] F. Liu, S. Shen, V. Anh and I. Turner, Analysis of a discrete non-Markovian random walk approximation for the time fractional diffusion equation, *ANZIAM J.*, 46(E) (2005), 488-504.
- [19] Ch. Lubich, Discretized fractional calculus [J], *SIAMJ. Math. Anal.*, vol. 17, no. 3, May (1986), pp. 704-719.
- [20] Mandelbrot, B., Some noises with  $1/f$  spectrum, a bridgre between direct current and white noise, *IEEE Transactions on In formation Theory* 13(2), (1967), 289-298.
- [21] R.J. Mark and M.W. Hall, Differentegral interpolation from a bandlimited signal's samples, *IEEE Trans. Acoust, Speech Signal Processing* 29 (1981) 872-877.
- [22] K. S. Miller, B. Boss, An introduction to the fractional calculus and fractional differential equations, Wiley, New Tork, USA, (1993).
- [23] K. B. Oldham, J. C. Spanier, The fractional calculus, Academic Press, San Diego, CA, USA, (1974).

- [24] I. Podlubny, Fractional differential equations, Academic Press, San Diego, CA, USA, (1999).
- [25] S. Shen and F. Liu, A computational effective method for fractional order Bagley-Torvik equation, Journal of Xiamen University (NATURAL Science), Vol.45, No.3, 2004, 306-311.
- [26] S. Shen and F. Liu, Error analysis of an explicit finite difference approximation for the space fractional diffusion equation with insulated ends, ANZIAM J., 46(E), (2005), 871-887.
- [27] S.B. Skaar, A.N. Michel and R.K. Miller, Stability of viscoelastic control systems, IEEE Transactions on a Automatic Control 3(4), (1988), 348-357.
- [28] H.H. Sun, N. Onaral and Y. Tsao, Application of position reality principle to metal electrode linear polarization phenomena, IEEE Transactions on Ciomedical Engineering 31(10), (1984), 664-674.
- [29] H.H. Sun, A.A. Abdelwahab and B. Onaral, Linear approximation of transfer function with a pole of fractional order, IEEE Transactions on Automatic Control 29(5), (1984), 441-444.
- [30] K. Diethelm and A.D. Freed, On the Solution of Nonlinear Fractional-Order Differential Equations Used in the Modeling of Viscoplasticity, Scientific Computing in Chemical Engineering II-Computational Fluid Dynamics, Reaction Engineering, and Molecular properties (1999), 217-224.
- [31] N.T. Shawagfeh, Analytical approximate solutions for nonlinear fractional differential equations, Applied Mathematics and Computation 131, (2002), 517-529.